



Extension

UNIVERSITY OF WISCONSIN-MADISON

Provided to you by:

University of Wisconsin Pest Alert

Lily Leaf Beetle

PJ Liesch, UW-Madison Entomology and Lisa Johnson, Extension Dane County

The lily leaf beetle (LLB), *Lilioceris lillii*, also known as the red or scarlet lily beetle, is an invasive insect of Eurasian origin. This insect was first reported in North America in eastern Canada during World War II and was most likely introduced in shipments of plant materials from Europe. LLB spread to New England in the 1990's and has been moving westward since that time. LLB made its first appearance in Wisconsin in 2014 and as of early 2023 has been found in 24 counties including Brown, Calumet, Clark, Dane, Door, Langlade, Lincoln, Marathon, Marquette, Milwaukee, Oneida, Outagamie, Pierce, Portage, Price, Shawano, Sheboygan, Taylor, Vernon, Vilas, Waukesha, Waushara, Waupaca and Wood Counties.



Severe feeding damage from lily leaf beetles on a lily plant (left) and an adult lily leaf beetle (right).
Photos courtesy of Leslie J. Mehrhoff, University of Connecticut, Bugwood.org

LLB can cause significant damage to true lilies (*Lilium* spp.), including both native and cultivated types, as well as fritillaries (*Fritillaria* spp.). LLB can also cause damage, but to a lesser extent, to lily of the valley (*Convallaria majalis*) and Solomon's seal (*Polygonatum* spp.). LLB does not cause damage to daylilies (*Hemerocallis* spp.), canna lilies (*Canna* spp.) or calla lilies (*Calla palustris*).

Appearance: LLB adults are bright red and approximately $\frac{1}{4}$ inch long, with black legs, head, antennae, and undersides. When disturbed, LLBs tumble from plants and land upside down on the ground, where they play dead. Their dark-colored undersides make them hard to spot. To potentially deter predators, adults squeak if squeezed.

LLB larvae have plump, squishy bodies and are slug-like in appearance. They tend to be orange or yellowish in color with black heads. At maturity, larvae can be almost $\frac{1}{2}$ inch long. As a defensive tactic, larvae typically camouflage themselves with their own excrement and can resemble slimy greenish-brown slugs or a mobile pile of animal droppings.

LLB eggs are tiny (less than $\frac{1}{10}$ inch long) and reddish in color. Female LLBs lay the eggs (typically in a batch of a dozen or less) on the underside of leaves in a row parallel to a vein.

Symptoms and Effects: Both LLB adults and larvae chew irregular holes and notches in lily leaves, stems, and developing buds. Larvae are the more damaging stage of the insect. When feeding damage is severe, LLBs can completely defoliate plants and destroy flowers.

Life Cycle: There is a single generation of LLBs each year, but adults can live for several years. LLBs overwinter as adults in sheltered places, soil, and plant debris in gardens and wooded areas.



Extension

UNIVERSITY OF WISCONSIN-MADISON

These overwintering sites are not necessarily near lilies or other host plants. Early in the spring, LLB adults emerge to feed, mate and lay eggs. Because they are strong fliers, LLBs can disperse over long distances to locate host plants. Eventually, females lay between 250 and 450 eggs. Eggs hatch within four to eight days. Emerging larvae feed for two to three weeks during the spring and early summer. Engorged larvae eventually drop to the soil to pupate (i.e., transform into adults). Pupae are bright orange and encased in a white cocoon with black spots. Adults emerge 16 to 22 days later and feed throughout the rest of the growing season and into the fall.



Lily leaf beetle eggs (left) and a lily leaf beetle larva camouflaged with its own feces. Photos courtesy of Gail Hampshire (left) and Kenneth R. Law, USDA APHIS PPQ, Bugwood.org

Control: If you have a small number of lilies, consider hand-picking and crushing adults and larvae or knocking them into a container of soapy water. Also, crush eggs by hand if you see them. Repeat this process regularly throughout the growing season. If LLB becomes a chronic and severe problem, consider replacing your lilies with plants that are not attacked by the insect.

You can also use conventional and organic insecticides to help protect plants from the LLB. Conventional insecticides containing carbaryl, cyfluthrin, cypermethrin, deltamethrin, lambda-cyhalothrin, permethrin, and zeta-cypermethrin control a broad range of pests, including LLB. Organic insecticides containing azadirachtin, pyrethrins, or spinosad can also be used, as well as horticultural oils and insecticidal soap. Make sure that the product that you select is labelled for use on landscape flowers. Be aware that conventional and organic insecticides can pose risks to pollinators, so follow all directions on the label to minimize risks to bees and other pollinators.

For more information on lily leaf beetle: Contact your county Extension agent.

© 2020-2021 by the Board of Regents of the University of Wisconsin System doing business as the Division of Cooperative Extension of the University of Wisconsin Extension.

An EEO/Affirmative Action employer, University of Wisconsin Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements. This document can be provided in an alternative format by calling Brian Hudelson at (608) 262-2863 (711 for Wisconsin Relay).

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Lisa Boyer, Julie Hill and Dan Marzu for reviewing this document.

A complete inventory of University of Wisconsin Garden Facts is available at the University of Wisconsin-Madison Division of Extension Plant Disease Diagnostics Clinic website: <https://pdcd.wisc.edu>.